

VEECH SURFACES AND COMPLETE PERIODICITY IN GENUS 2

KARIANE CALTA

ABSTRACT. We announce several results pertaining to Veech surfaces and completely periodic translation surfaces in genus 2. A translation surface is a pair (M, ω) where M is a Riemann surface and ω is an abelian differential on M . Equivalently, a translation surface is a two-manifold which has transition functions which are translations and a finite number of conical singularities arising from the zeroes of ω . A saddle connection is a geodesic segment connecting two (not necessarily distinct) conical singularities with no singularities in its interior.

A direction v on a translation surface is completely periodic if any trajectory in the direction v is either closed or ends in a singularity, i.e. if the surface decomposes as a union of cylinders in the direction v . Then, we say that a translation surface is completely periodic if any direction in which there is a saddle connection joining a singularity to itself or at least one cylinder of closed trajectories is completely periodic. There is an action of the group $SL(2, \mathbb{R})$ on the space of translation surfaces. A surface which has a lattice stabilizer under this action is said to be Veech. Veech proved that any Veech surface is completely periodic, but the converse is false.

In this announcement, we use the " J -invariant" of Kenyon and Smillie to obtain a classification of all Veech surfaces in the space $\mathcal{H}(2)$ of genus 2 translation surfaces with corresponding abelian differentials which have a single double zero. Furthermore, we obtain a classification of all completely periodic surfaces in genus 2.

A translation surface is a pair (M, ω) where M is a closed Riemann surface and ω is an abelian differential on M . Away from the zeroes of ω , a chart z can be chosen so that $\omega = dz$, which determines a Euclidean metric in that chart. The change of coordinates away from the zeroes of ω are of the form $z \rightarrow z + c$. In the neighborhood of a zero, one can choose a coordinate z so that $\omega = z^k dz$. The total angle about this zero is $2(k+1)\pi$ and the order of the zero is k . Such zero is called a conical singularity. As a result of this definition, a translation surface can also be thought of as a two-manifold with transition functions which are translations and which has a finite number of conical singularities, arising from the zeroes of ω . Each of these singularities has total angle $2\pi n$ for $n \in \mathbb{Z}^+$. Thus, one can also think of a translation surface as a finite number of polygons in the plane $\mathbb{R}^2$, glued along parallel sides. Throughout, we will often denote a translation surface by S , and this notation should be thought of as incorporating both the Riemann surface M and the differential ω .

If S is a translation surface, a *saddle connection* is a geodesic connecting two (not necessarily distinct) conical singularities of S which has no singularities in its interior. Note that since the metric on S is Euclidean away from the zeroes of ω , saddle connections are straight lines, but in general, geodesics are unions of saddle connections where the line segments comprising the geodesic are allowed to switch directions at the conical singularities of S . A closed geodesic on S determines a cylinder of parallel freely homotopic closed curves such that the boundaries of the cylinder consist of unions of saddle connections. A direction v on S is said to be *completely periodic* if all the trajectories in the direction v are either closed or terminate at a singularity. In other words, v is completely periodic if in the direction v ,

the surface decomposes as a union of cylinders. A surface S is *completely periodic* if any direction in which there is at least one cylinder or a saddle connection connecting a zero to itself is completely periodic.

It can be shown that if a translation surface S has genus $g \geq 1$, then the number of zeroes of ω , counting multiplicity, is $2g - 2$. Let α be a partition of $2g - 2$, that is a collection of positive integers $\alpha_1, \dots, \alpha_k$ such that $\alpha_1 + \dots + \alpha_k = 2g - 2$. Thus, if ω has k distinct zeroes, and α_i is the multiplicity of one of the k zeroes, then ω determines a partition of $2g - 2$. Now, fix a genus g and a partition α of $2g - 2$. We let $\mathcal{H}(\alpha)$ denote the space of translation surfaces (M, ω) where M is a Riemann surface of genus g and ω is an abelian differential on M whose zeroes correspond in the way described above to the partition α . For example, the set of translation surfaces of genus 2 is stratified as $\mathcal{H}(1, 1) \cup \mathcal{H}(2)$, where in $\mathcal{H}(1, 1)$ the 1-form has two distinct simple zeroes, and in $\mathcal{H}(2)$ it has a single double zero.

There is an action of the group $SL(2, \mathbb{R})$ on each stratum $\mathcal{H}(\alpha)$. If we think of a translation surface S as a two-manifold with transition functions which are translations and which has a finite number of conical singularities, and if $g \in SL(2, \mathbb{R})$, then gS is the two manifold which has charts which are given by $g \circ f_i$ where f_i is a chart of S . It is easily checked that the transition functions of gS are still translations and the number of singularities and their total angles are preserved. Alternatively, if $S = P_1 \cup \dots \cup P_n$, where each P_i is a polygon in the plane $\mathbb{R}^2$, then $gS = gP_1 \cup \dots \cup gP_n$. A translation surface is said to be *Veech* if its stabilizer under this action of $SL(2, \mathbb{R})$ is a lattice. It is a theorem of Veech that Veech surfaces are completely periodic, but the converse is not true.

For a curve $\gamma \subset S$, let $p(\gamma)$ denote $\int_\gamma(\omega)$, where ω is the 1-form defining the translation surface structure. We identify $\mathbb{R}^2$ with $\mathbb{C}$. Define a translation surface S to be *quadratic* if there exists $d > 0$ (not a perfect square) such that $p(H_1(S, \mathbb{Z})) \subset \mathbb{Q}(\sqrt{d}) \times \mathbb{Q}(\sqrt{d})$. Any Veech surface in genus 2 (perhaps after rescaling) is known to be quadratic.

Let C be a cylinder of periodic trajectories in the direction v . We use the term *width* for the length of the periodic trajectory, and *height* for the distance across the cylinder in the orthogonal direction. If w is any other direction, we can talk about the *twist* of C along w . Let $v^\perp$ be the direction orthogonal to v given by the right hand rule, and let w' be the vector in the direction w whose projection onto $v^\perp$ has length equal to the height of the cylinder. Now, suppose that the angle θ between v and w satisfies $0 \leq \theta \leq \pi/2$. Then we can define the twist t in the direction w to be the length of the projection of w' onto v modulo the width of the cylinder. If $\pi/2 < \theta \leq \pi$, then we define the twist to be the width of the cylinder minus the length of the projection of w' onto $-v$ modulo the width of the cylinder.

Theorem 1. *Let S be a Veech surface in $\mathcal{H}(2)$ which cannot be rescaled so that $p(H_1(S, \mathbb{Z})) \subset \mathbb{Q} \times \mathbb{Q}$. Let v be a completely periodic direction for S . Suppose the cylinder decomposition in the direction v has two cylinders. Let $w_1, w_2, h_1, h_2, t_1, t_2$ be the widths, heights and twists of these two cylinders. The twists are measured along some direction w which we assume is also completely periodic. After rescaling the surface, we may assume that these variables are algebraic integers in $\mathbb{Q}(\sqrt{d})$ where d is some square-free, positive integer. Then the following equations are satisfied:*

$$(1) \quad w_1 \bar{h}_1 = -w_2 \bar{h}_2$$

$$(2) \quad \bar{w}_1 t_1 + \bar{w}_2 t_2 + \bar{w}_1 w_2 = w_1 \bar{t}_1 + w_2 \bar{t}_2 + w_1 \bar{w}_2, \quad 0 \leq t_1 < w_1, 0 \leq t_2 < w_2$$

(where the bar denotes conjugation in $\mathbb{Q}(\sqrt{d})$.)

Conversely, let S be a surface in $\mathcal{H}(2)$. Suppose there exists a direction v in which there is a cylinder decomposition for which the heights, widths and twists (along some direction w) belong to $\mathbb{Q}(\sqrt{d})$ and satisfy (1) and (2). Then S is Veech.

(If $S \in \mathcal{H}(2)$ is Veech and has a one cylinder decomposition in some direction v , then $p(H_1(S, \mathbb{Z})) \subset \mathbb{Q} \times \mathbb{Q}$.)

We note that a different classification of Veech surfaces in genus 2, formulated in terms of Jacobians with real multiplication, was obtained by C. McMullen in [M].

Let $\mathcal{O}_d^+$ denote the positive algebraic integers in $\mathbb{Q}(\sqrt{d})$.

Lemma 1. *For fixed integers c_1 and c_2 , the equations (1), (2) and*

$$(3) \quad w_1 h_1 + w_2 h_2 = 2(c_1 + c_2 \sqrt{d})$$

have finitely many solutions in $\mathcal{O}_d^+$, up to the action by the group of units. Let us denote this finite number by $H(c_1, c_2)$.

Let Ω be a set containing one element for each Veech surface in $\mathcal{H}(2)$, where we identify any two surfaces in the same $GL(2, \mathbb{R})$ orbit. Let P denote the set of positive solutions in algebraic integers in $\mathbb{Q}(\sqrt{d})$ to (1) and (2). Then in view of Theorem (1) there exists a map $f_1 : P \rightarrow \Omega$. Now if S is a Veech surface, then there is a surface S' in the $GL(2, \mathbb{R})$ orbit of S such that $p(H_1(S'))$ is a primitive sublattice of $\mathcal{O}_d \times \mathcal{O}_d$ with no common factor. The area of S' is independent of the choice of S' . Let $f_2 : \Omega \rightarrow \mathcal{O}_d^+$ denote the map which sends (the equivalence class of) S to the area of S' .

Corollary 1. (Classification of Veech surfaces in $\mathcal{H}(2)$). *The maps $f_1 : P \rightarrow \Omega$ and $f_2 : \Omega \rightarrow \mathcal{O}_d^+$ are finite-to-one. In fact, the cardinality of $(f_1 \circ f_2)^{-1}(2(c_1 + c_2 \sqrt{d}))$ is $H(c_1, c_2)$.*

We now state some results for $\mathcal{H}(1, 1)$. Suppose v is a completely periodic direction for $S \in \mathcal{H}(1, 1)$. Without loss of generality, we may assume that in the direction v , S decomposes into three cylinders, and for $1 \leq i \leq 3$, let w_i , h_i and t_i denote the widths, heights and twists. After renumbering, we may assume that $w_3 = w_1 + w_2$. Denote $s_1 = h_1 + h_3$, $s_2 = h_2 + h_3$, $\tau_1 = t_1 + t_3$, $\tau_2 = t_2 + t_3$. For $c_1 + c_2 \sqrt{d} \in \mathcal{O}_d^+$, consider the following equations:

$$(4) \quad w_1 \bar{s}_1 = -w_2 \bar{s}_2$$

$$(5) \quad \bar{w}_1 \tau_1 + \bar{w}_2 \tau_2 + \bar{w}_1 w_2 = w_1 \bar{\tau}_1 + w_2 \bar{\tau}_2 + w_1 \bar{w}_2, \quad 0 \leq t_1 < w_1, 0 \leq t_2 < w_2$$

$$(6) \quad w_1 s_1 + w_2 s_2 = 2(c_1 + c_2 \sqrt{d})$$

Theorem 2. *Let $\mathcal{M}(c_1, c_2)$ denote the set of surfaces in $\mathcal{H}(1, 1)$ which are completely periodic and for each completely periodic direction v , the lengths heights and twists of the cylinder decomposition in that direction satisfy (4), (5) and (6). Then $\mathcal{M}(c_1, c_2)$ is a closed subset of dimension 3 which is invariant under the $SL(2, \mathbb{R})$ action. It is non-empty provided (4) (5) and (6) have a solution in $\mathcal{O}_d^+$, and any Veech surface in $\mathcal{H}(1, 1)$ is contained in such a subset.*

We also obtain a classification of the completely periodic surfaces in genus 2. In fact

Theorem 3. *In $\mathcal{H}(2)$ every completely periodic surface is Veech. In $\mathcal{H}(1, 1)$, a surface is completely periodic if it has a cylinder decomposition in some direction v , such that the heights widths and twists satisfy (4) (5) and (6). Conversely, if $S \in \mathcal{H}(1, 1)$ is completely*

periodic then after rescaling, either $p(H_1(S)) \subset \mathbb{Q} \times \mathbb{Q}$ or $p(H_1(S)) \subset \mathbb{Q}(\sqrt{d}) \times \mathbb{Q}(\sqrt{d})$ for some square-free $d > 0$. If S cannot be rescaled so that $p(H_1(S)) \subset \mathbb{Q} \times \mathbb{Q}$ and if v is any completely periodic direction, then (4) (5) and (6) hold, where the w_i, h_i, t_i denote the widths, heights and twists of the cylinder decomposition along v (and the twists are measured along another completely periodic direction v').

The proofs of the results stated here use the J invariant for translation surfaces defined by Kenyon and Smillie in their paper “Billiards on rational angled triangles” as follows. First if P is a polygon in $\mathbb{R}^2$ with vertices $v_1, \dots, v_n$ in counterclockwise order about the boundary of P , then $J(P) = v_1 \wedge v_2 + \dots + v_{n-1} \wedge v_n + v_n \wedge v_1$. Then, if S is a translation surface with a cellular decomposition into planar polygons $P_1 \cup \dots \cup P_k$, $J(S) = \sum_{i=1}^k J(P_i)$. Furthermore, there are two linear projections J_{xx} and J_{yy} from $\mathbb{R}^2 \wedge_{\mathbb{Q}} \mathbb{R}^2$ to $\mathbb{R} \wedge_{\mathbb{Q}} \mathbb{R}$ given on basis elements as

$$\begin{aligned} J_{xx} \left(\begin{pmatrix} a \\ b \end{pmatrix} \wedge \begin{pmatrix} c \\ d \end{pmatrix} \right) &= a \wedge c \\ J_{yy} \left(\begin{pmatrix} a \\ b \end{pmatrix} \wedge \begin{pmatrix} c \\ d \end{pmatrix} \right) &= b \wedge d \end{aligned}$$

and one linear projection J_{xy} from $\mathbb{R}^2 \wedge_{\mathbb{Q}} \mathbb{R}^2$ to $\mathbb{R} \otimes_{\mathbb{Q}} \mathbb{R}$ defined on basis elements as

$$J_{xy} \left(\begin{pmatrix} a \\ b \end{pmatrix} \wedge \begin{pmatrix} c \\ d \end{pmatrix} \right) = a \otimes d - c \otimes b.$$

Let v be some direction on a translation surface S . If v is either the horizontal or vertical direction, J_{vv} has already been defined. Suppose that v is neither the horizontal nor vertical direction, and let the vector $\begin{pmatrix} 1 \\ q \end{pmatrix}$ represent the direction v . If g is any matrix in $SL(2, \mathbb{R})$ such that $g\begin{pmatrix} 1 \\ q \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, then we define $J_{vv}(S) = J_{yy}(gS)$. It can be shown that this quantity is independent of the choice of matrix g . Also, given a pair of linearly independent directions v, w on S , let v', w' be vectors in the directions v and w so that there exists an element $g \in SL(2, \mathbb{R})$ such that $gv' = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $gw' = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ and define $J_{vw}(S) = J_{xy}(gS)$. (In the case that $v = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $w = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$, define $J_{vw}(S) = J_{yx}(S)$ where $J_{yx}(\begin{pmatrix} a \\ b \end{pmatrix} \wedge \begin{pmatrix} c \\ d \end{pmatrix}) = b \otimes c - d \otimes a$.) It can be shown that a different choice of v', w' will multiply each of the components of each of the tensors comprising J_{vw} by the same number.

An alternative formula for the Kenyon-Smillie invariant is:

$$J = 2 \sum_{i=1}^g p(a_i) \wedge p(b_i)$$

where a_i, b_i is a symplectic homology basis for the surface S .

It is proven in Kenyon and Smillie that if the horizontal direction x on a translation surface S is completely periodic, then $J_{yy} = 0$.

We define a direction v to be *homological* if there exists $\lambda \in H_1(S)$ and $r \in \mathbb{R}$ such that $rv = p(\lambda)$. Note that λ is not assumed to be simple.

We say that a translation surface S has **Property X** if for any homological direction v on S , $J_{vv} = 0$. I have proven the following theorems regarding property X.

Theorem 4. *If a genus 2 translation surface has property X, then it is completely periodic.*

Theorem 5. *If a translation surface is completely periodic, then it has property X. In particular, any Veech surface has Property X.*

Theorem 6. *If a surface in $\mathcal{H}(2)$ has property X, then it is Veech.*

Proposition 1. *Let S be a quadratic translation surface which cannot be rescaled so that $p(H_1(S, \mathbb{Z})) \subset \mathbb{Q} \times \mathbb{Q}$. Let v and w be any two linearly independent directions such that $J_{vv} = J_{ww} = 0$. We may write $J_{vw}(S) = c_1(1 \otimes 1) + c_2(1 \otimes \sqrt{d}) + c_3(\sqrt{d} \otimes 1) + c_4(\sqrt{d} \otimes \sqrt{d})$ where $c_i \in \mathbb{Q}$. Then S has property X if and only if $c_2 = c_3$ and $c_1 = dc_4$.*

A straightforward calculation shows the following:

Lemma 2. *Let $S \subset \mathcal{H}(2)$ be a quadratic translation surface which cannot be rescaled so that $p(H_1(S, \mathbb{Z})) \subset \mathbb{Q} \times \mathbb{Q}$. Assume that S has a two cylinder decomposition in the horizontal direction. Let $w_1, w_2, h_1, h_2, t_1, t_2$ be the widths, heights and twists of these two cylinders. Then $J_{xx} = 0$ if and only if (2) holds. Furthermore the equations $c_1 = dc_4$ and $c_2 = c_3$ defining property X are equivalent to (1).*

Now Theorem 1 follows from Theorem 5, Theorem 6, Proposition 1, and Lemma 2. The proof of Theorem 6 uses Theorem 4 and Lemma 1. The proof of Theorem 2 also uses Theorem 4 and Lemma 1.

We say that a translation surface S is **hyperperiodic** if any homological direction on S is completely periodic. Clearly hyperperiodicity implies both complete periodicity and property X. The following theorem was obtained in joint work with A. Eskin and M. Boshernitzan (see [BCE]).

Theorem 7. *Any quadratic Veech surface is hyperperiodic. Also, in genus 2, hyperperiodicity, complete periodicity and property X are equivalent.*

We note that Theorem 7 gives a complete description of the periodic directions in e.g. the regular pentagon. We also note that hyperperiodicity remains invariant under the deformation where the absolute homology is fixed and the relative homology changes.

REFERENCES

- [M] C. McMullen. Billiards and Teichmüller curves on Hilbert modular surfaces. *Preprint*.
- [BCE] M. Boshernitzan, K. Calta, A. Eskin. *In preparation*.

Department of Mathematics, University of Chicago, Chicago IL 60637, USA;
kcalta@math.uchicago.edu